


ADA 081960

12



LEVEL II

This document has been approved  
for public release and sale; its  
distribution is unlimited.

The Florida State University  
Department  
of  
Statistics  
Tallahassee, Florida  
32306



12

6

THE EXTREMAL LARGE DEVIATION RATE  
OF THE F-STATISTIC

10

L. / Barker J. / Sethuraman

9

FSU Statistics Report M529  
Technical Report No. D-42

11

October 1979

The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306

DTIC  
SELECTED  
D

12) 19

15) DAA G 29-76-G-0278

18) ARO

19) 16713.7-M

Research partially supported by United States Army Research Office Grant  
number DAA29-76-G00278.

14) FSU-STATISTICS-M529, TR-D-42-ARO

The contents of this report have been reviewed for technical accuracy and approved for publication, but they do not represent an official Army position or that of the originating organization.

This document has been approved  
for public release and sale; its  
distribution is unlimited.

44714

# ABSTRACT

→ The extremal large deviation rate of the F-statistic over a large class is determined. It is shown that the Bahadur efficiency of the F-test relative to Moses' rank test for equality of variances is zero at every alternative when the distribution of the underlying observations is unspecified. ↗

Key words: Bahadur efficiency, extremal large deviations, F-statistic, nonparametric tests.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DOC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
A	

Page - A -

I. Introduction. Extremal large deviation rates measure the slowest rate of convergence to zero of certain sequences of probabilities. When these probabilities arise from a sequence of test statistics, extremal large deviation rates find application in the computation of asymptotic relative efficiency (ARE) of tests, such as Bahadur's definition of ARE.

The extremal large deviation rate of only a few common statistics is known. Here we will determine the extremal large deviation rate of the F-statistic.

Before precisely defining extremal large deviation rates, we will discuss Bahadur's definition of ARE, in which it finds application. Let  $(X, A)$  be a measurable space. Let  $X = \{X_i\}_{i=1}^{\infty}$  be a sequence of random variables on  $(X, A)$ , and let  $P$  be a probability measure on  $(X, A)$  such that  $X$  is a sequence of independent identically distributed random variables with common distribution  $P_P$ . Let  $C$  and  $C'$  be disjoint non-empty collections of distributions, and consider the problem of testing

$$H_0 : P \in C \text{ vs. } H_a : P \in C'.$$

Typically, decision rules are of the form "reject  $H_0$  if  $S_n$  exceeds some specified value  $a$ ", where  $S_n$  is a measurable real valued function of  $\{X_i\}_{i=1}^n$ ; that is  $S = \{S_i\}_{i=1}^{\infty}$  is a sequence of test statistics. Call this test  $T_S(a)$ .

Let  $S' = \{S'_i\}_{i=1}^{\infty}$  be another sequence of test statistics. Call the test that rejects  $H_0$  when  $S'_n$  exceeds a specified value, say  $a'$ ,  $T_{S'}(a')$ . The test  $T_{S'}(a')$  is a competitor to the test  $T_S(a)$ . We choose between these tests by considering their ARE.

The most important definitions of ARE are due to Pitman, (1949), Hodges-Lehmann (1956), and Bahadur (1960). Under mild regularity conditions, evaluation

Acquisition For	NTIS	Card	✓
DOC TAB	Unannounced	Justification	
By	Distribution	Availability	
Dist	Availability	Special	

of the Bahadur efficiency of  $T_S(a)$  relative to  $T_{S'}(a')$  requires the determination of extremal large deviations rates of  $\{S_i\}_{i=1}^{\infty}$  and  $\{S'_i\}_{i=1}^{\infty}$ .

We now make precise the concept of an extremal large deviation rate.

Definition 1: If

$$W(S, C, a) = -\lim_{n \rightarrow \infty} n^{-1} \log \sup_{P \in C} P(S_n \geq a)$$

is well defined, we say that  $W(S, C, a)$  is the extremal large deviation rate at a deviation of a for the sequence  $S$  under the class of probability models  $C$ .

That is,  $W(S, C, a)$  is a measure of the rate at which

$$\sup_{P \in C} P(S_n \geq a)$$

converges to zero.

Let  $C$  be such that for every  $P \in C$ ,

$$P(S_n \rightarrow 0) = 1.$$

Consider the test  $T_S(a)$ , where  $a > 0$ . Now  $F_n^*(a) = \sup_{P \in C} P(S_n \geq a)$  is the significance level of  $T_S(a)$ , and  $L_n = F_n^*(S_n)$  is the usual observed significance level. If  $P \in C$ ,  $P(L_n \rightarrow 0) = 0$ . If  $P \notin C$  (in particular, if  $P \in C'$ , the alternative class), then  $P(L_n \rightarrow 0) = 1$ . In typical cases,  $L_n$  converges to zero exponentially fast at non-null  $P$ .

Definition 2: The half slope of  $T_S(\cdot)$  at non-null  $P$  is given by

$$r(P) = -\lim_{n \rightarrow \infty} n^{-1} \log L_n [P]$$

when the above is well defined.

We may interpret  $\rho(P)$  as a measure of the rate at which the observed significance level tends to zero, with large values indicating more rapid convergence. Bahadur has exploited this property to define a measure of ARE, described below.

Definition 3: Let  $T_S(\cdot)$  and  $T_{S'}(\cdot)$  be as defined earlier. At non-null  $P$ , let  $T_S(\cdot)$  have half slope  $\rho(P)$  and let  $T_{S'}(\cdot)$  have half slope  $\rho'(P)$ . The Bahadur efficiency of  $T_S(\cdot)$ , relative to  $T_{S'}(\cdot)$ , at non-null  $P$  is given by  $\rho(P)/\rho'(P)$ .

The following theorem, due to Bahadur, relates the concepts of half slope and extremal large deviation rates.

Theorem 1: (Bahadur, 1960). Let  $P(S_n \rightarrow a(P)) = 1$  at non-null  $P$ . Let  $W(S, C, t)$  be well defined for  $t$  in a neighborhood of  $a(P)$  and continuous at  $t = a(P)$ . Then  $\rho(P) = W(S, C, a(P))$ .

Consider the case  $S_n = n^{-1} \sum_{i=1}^n X_i$ , where  $C$  consists of a single distribution  $F$  and  $\int x dF(x) = 0$ . Using Chernoff's Theorem, stated below, evaluation of the half slope of  $\bar{S}$  at a fixed non-null distribution is straightforward.

Theorem 2: (Chernoff, 1952). Let  $\{X_i\}_{i=1}^{\infty}$  be a sequence of i.i.d. random variables. Let  $\phi(t) = E e^{tx_1}$ , not necessarily finite for  $t \neq 0$ . Then for every real  $a$ ,

$$\lim_{n \rightarrow \infty} n^{-1} \log P(n^{-1} \sum_{i=1}^n X_i \geq a) = \inf_{t \geq 0} \log \phi(t) e^{-at}.$$

Hence, at non-null  $G$ , the half slope of  $\bar{S}$  equals  $-\inf_{t \geq 0} \log \psi(t) e^{-at}$  where  $\psi(t) = \int e^{tx} dF(x)$  and  $a = \int x dG(x)$ .

## II. Large Deviation Rates of the F-Statistic

Extremal large deviation rates over a large class have only been evaluated in a few instances. For example, if the statistic of interest is a linear rank statistic, a straightforward method exists. See Klotz (1965), who evaluates the half slope of the Wilcoxon rank sum statistic, as well as other linear rank statistics.

One of the few non-distribution-free statistics for which the extremal large deviation rate has been studied is the t-statistic.

Theorem 3: (Jones and Sethuraman, 1978). Let  $\{X_i\}_{i=1}^{\infty}$ ,  $\{Y_i\}_{i=1}^{\infty}$  be two sequences of mutually independent identically distributed random variables, with common distribution F and joint distribution  $P_F$ . Let

$$\bar{X}_n = n^{-1} \sum_{i=1}^n X_i, \quad \bar{Y}_n = n^{-1} \sum_{i=1}^n Y_i,$$

$$s_{x,n}^2 = n^{-1} \sum_{i=1}^n X_i^2 - (\bar{X}_n)^2, \text{ and}$$

$$s_{y,n}^2 = n^{-1} \sum_{i=1}^n Y_i^2 - (\bar{Y}_n)^2.$$

Define the one- and two-sample t-statistics,  $T_{1,n}$  and  $T_{2,n}$  respectively, as

$$T_{1,n} = \bar{X}_n / (s_{x,n}^2)^{1/2} \text{ and}$$

$$T_{2,n} = (\bar{X}_n - \bar{Y}_n) / ((s_{x,n}^2 + s_{y,n}^2)/2)^{1/2}$$

where expressions of the form  $r/0$  are interpreted as

$$\begin{cases} +\infty & \text{if } r > 0 \\ 0 & \text{if } r = 0 \\ -\infty & \text{if } r < 0 \end{cases}.$$

Let  $C_1 = \{F: F \text{ is symmetric about } 0\}$  and let  $C_2 = \{(F, G): F = G\}$ . Then for every  $a > 0$ ,  $j = 1$  or  $2$ ,  $W(\{T_{j,n}\}_{n=1}^{\infty}, C_j, ja) = jI(\frac{1}{2}(1 + a/(1 + a^2)^{\frac{1}{2}}), \frac{1}{2})$ , where for  $0 < p_1, p_2 < 1$ ,  $I(p_1, p_2) = p_1 \log \frac{p_1}{p_2} + (1 - p_1) \log \frac{1 - p_1}{1 - p_2}$ . Further, this rate of convergence is achieved by a t-statistic based on symmetric Bernoulli observations. That is, the t-statistic converges most slowly when based on symmetric Bernoulli observations.

Definition 4: Let  $\{X_i\}_{i=1}^{\infty}$ ,  $\{Y_i\}_{i=1}^{\infty}$ ,  $s_{x,n}^2$ ,  $s_{y,n}^2$ , and indeterminate forms be as in Theorem 3. Let the F-statistic,  $F(n)$ , be given by  $s_{x,n}^2/s_{y,n}^2$ . Let the shifted F-statistic,  $F_s(n)$ , equal  $F(n) - 1$ .

It is reasonable to base a test of equality of variances, assumed to exist, on  $F_s(n)$ , with large values indicating rejection. Under the null hypothesis,  $F_s(n) \rightarrow 0$  a.s. At an alternative  $H_a$ ,  $F_s(n) \rightarrow a(H_a)$  a.s., where  $a(H_a)$  is assumed positive. We will now determine the large deviation rate of  $F_s(n)$  under certain parametric assumptions.

Theorem 4: Let  $F$  assign mass  $\frac{1}{2}$  each to  $\pm 1$ . That is,  $F$  is a symmetric Bernoulli distribution. Let  $\{X_i\}_{i=1}^{\infty}$ ,  $\{Y_i\}_{i=1}^{\infty}$  be mutually independent identically distributed random variables, with common distribution  $F$ . Let  $F_s(n)$  be based on  $\{X_i\}_{i=1}^{\infty}$ ,  $\{Y_i\}_{i=1}^{\infty}$ . Then for any  $a > 0$ ,

$$-\lim_{n \rightarrow \infty} n^{-1} \log P(F_s(n) \geq a) = I(\frac{1}{2}(1 + (a/1 + a)^{\frac{1}{2}}), \frac{1}{2})$$

where  $I(\cdot, \cdot)$  is as in Theorem 3.

Proof: For any real  $a$ ,

$$P(F_s(n) \geq a) = P(s_{x,n}^2/s_{y,n}^2 \geq 1 + a) = P((1 - \bar{X}_n^2)/(1 - \bar{Y}_n^2) \geq 1 + a) =$$

$$P((1 + a)\bar{Y}_n^2 - \bar{X}_n^2 \geq a) \leq P(\bar{Y}_n^2 \geq a/(1 + a)).$$



By Chernoff's Theorem,

$$1 \quad \lim_{n \rightarrow \infty} n^{-1} \log P(Y_n^2 \geq a/(1+a)) = (-I(\frac{1}{2}(1 + (a/a+1)^{\frac{1}{2}}), \frac{1}{2})).$$

Choose and fix  $\epsilon$ ,  $1 > \epsilon > 0$ . Now  $P(F_S(n) \geq a) = P((1 - \bar{X}_n^2)/(1 - \bar{Y}_n^2) \geq 1+a) \geq P(\bar{Y}_n^2 \geq (a+\epsilon)/(a+1), \bar{X}_n^2 < \epsilon) = P(\bar{Y}_n^2 \geq (a+\epsilon)/(a+1)) P(\bar{X}_n^2 < \epsilon)$ .

But  $\lim_{n \rightarrow \infty} \log P(\bar{X}_n^2 < \epsilon) = 0$ ;

$$\begin{aligned} \text{Hence } \lim_{n \rightarrow \infty} \lim_{n \rightarrow \infty} n^{-1} \log P(F_S(n) \geq a) &\geq \\ \lim_{n \rightarrow 0} \lim_{n \rightarrow \infty} n^{-1} \log P(\bar{Y}_n^2 \geq (a+\epsilon)/(a+1)) &= \\ \lim_{\epsilon \rightarrow 0} -I(\frac{1}{2}(1 + ((a+\epsilon)/(a+1))^{\frac{1}{2}}), \frac{1}{2}) &= \\ I(\frac{1}{2}(1 + ((a+\epsilon)/(a+1))^{\frac{1}{2}}), \frac{1}{2}). \end{aligned}$$

By lines 1 and 2,

$$-\lim_{n \rightarrow \infty} n^{-1} \log P(F_S(n) \geq a) = I(\frac{1}{2}(1 + (a/a+1)^{\frac{1}{2}}), \frac{1}{2}). \quad \square$$

**Theorem 5:** Let  $F$  be a normal distribution, with mean  $\mu$  and variance  $\sigma^2 > 0$ .

Let  $\{X_i\}_{i=1}^{\infty}, \{Y_i\}_{i=1}^{\infty}$  be mutually independent identically distributed random variables with common distribution  $F$ . Let  $F_S(n)$  be based on  $\{X_i\}_{i=1}^n, \{Y_i\}_{i=1}^n$ .

Then for any  $a \geq 0$ ,

$$-\lim_{n \rightarrow \infty} n^{-1} \log P(F_S(n) \geq a) = \log(2(a+1)^{\frac{1}{2}}/(a+2)).$$

**Proof:** Without loss of generality, let  $\mu = 0, \sigma^2 = 1$ . Let  $\{U_i\}_{i=1}^{\infty}, \{V_i\}_{i=1}^{\infty}$  be two sequences of mutually independent identically distributed random variable with common  $\chi_1^2$  distribution. Then for every  $a \geq 0$ ,

$$\begin{aligned} P(F_S(n) \geq a) &= \\ P(\sum_{i=1}^{n-1} U_i / \sum_{i=1}^{n-1} V_i \geq a+1) &= \\ P(\sum_{i=1}^{n-1} U_i \geq \sum_{i=1}^{n-1} (a+1)V_i) &= \\ P(\sum_{i=1}^{n-1} (U_i - (a+1)V_i) / (n-1) \geq 0). \end{aligned}$$

By Chernoff's Theorem

$$\lim_{n \rightarrow \infty} n^{-1} \log P\left(\sum_{i=1}^{n-1} (U_i - (a+1)V_i)/(n-1) \geq 0\right) =$$

$$\inf_{t \geq 0} \log(1 + 2at - 4(a+1)t^2)^{-1/2} =$$

$$\log(2(a+1)^{1/2}/(a+2)). \quad \square$$

In Table 1, we display  $I(\frac{1}{2}(1 + (a/(a+1))^{1/2}), \frac{1}{2})$  and  $-\log(2(a+1)^{1/2}/(a+2))$  for selected values of  $a > 0$ . We see that  $F_S(n)$  converges to zero more rapidly under some alternatives when the null distribution of the underlying observations is normal than if the null distribution of the underlying observations is symmetric Bernoulli. Unlike the t-statistic, the F-statistic does not converge most slowly when the underlying observations have a symmetric Bernoulli distribution.

III. Extremal Large Deviation Rates of the F-statistic. Before dealing with the F-statistic, which is used to test for equality of variances, we will consider a related statistic used for testing the equality of means of positive random variables.

Definition 4: Let  $\{X_i\}_{i=1}^{\infty}, \{Y_i\}_{i=1}^{\infty}$  be mutually independent identically distributed random variables with common distribution  $F \in G$ , where

$G = \{F: \int_0^{\infty} dF = 1, 0 < \int_0^{\infty} x dF(x) < \infty\}$ . Define the ratio statistic,  $R_n$ , as  $(\bar{X}_n / \bar{Y}_n) - 1$ , where  $\bar{X}_n, \bar{Y}_n$ , and indeterminate forms are as in Theorem 3.

The statistic  $R_n$  can be used to test  $H_0: EX_1 = EY_1$ . Under  $H_0$ ,  $R_n \rightarrow 0$  a.s. Under a non-null  $F$ ,  $R_n \rightarrow a(F)$  a.s. Without loss of generality, assume  $a(F) > 0$ .

Theorem 5. Let  $\{X_i\}_{i=1}^{\infty}, \{Y_i\}_{i=1}^{\infty} \in G$ , and  $R_n$  be as in Definition 4. Then for every  $a > 0$ ,

$$W(R_n, G, a) = 0.$$

Proof: Let  $F$  be a fixed but unspecified distribution in  $G$ . Clearly,

$$0 \leq W(R_n, G, a) \leq -\lim_{n \rightarrow \infty} n^{-1} \log P_F(R_n \geq a).$$

Let  $\epsilon > 0$  be chosen and fixed. Let  $\alpha = \epsilon / \log(1 + (a^2)/4(a+1))$ , and let  $\Gamma$  be a Gamma  $(\alpha, 1)$  distribution. Then

$$\lim_{n \rightarrow \infty} n^{-1} \log P_F(R_n \geq a) = \lim_{n \rightarrow \infty} n^{-1} \log P_F\left(\sum_{i=1}^n X_i \geq (a+1) \sum_{i=1}^n Y_i\right) =$$

$$\lim_{n \rightarrow \infty} n^{-1} \log P_F\left(\sum_{i=1}^n X_i - (a+1)Y_i \geq 0\right) = \log \inf_{t \geq 0} \psi(t) \psi(-(a+1)t),$$

by Chernoff's Theorem, where  $\psi(t) = (1-t)^{-\alpha}$ .

By elementary methods,  $\log \inf_{t \geq 0} \psi(t) \psi(-(a+1)t) = -\epsilon$ . The result follows, taking the limit as  $\epsilon \rightarrow 0$ .  $\square$

Before determining the extremal large deviation rate of the F-statistic, some preliminary results are needed.

Lemma 1. Let  $x$  tend to zero from above. Then  $I(\frac{1}{2}(1+x), \frac{1}{2}) = x^2/2 + o(x^2)$ .

Proof: Expanding  $I(\frac{1}{2}(1+x), \frac{1}{2})$  as a Taylor series, the result follows.

Lemma 2 (Jones and Sethuraman, 1978). Let  $\{X_i\}_{i=1}^{\infty}$  be a sequence of i.i.d. symmetric random variables.

Then for every sample size  $n$  and  $b \in (0, 1)$

$$P\left(\sum_{i=1}^n X_i / \left(\sum_{i=1}^n X_i^2\right)^{1/2} \geq n^{1/2} b\right) \leq \exp(-nI(\frac{1}{2}(1+b), \frac{1}{2})).$$

We are now ready to determine the extremal large deviation rate of the F-statistic over appropriate classes.

Theorem 7. Let  $F = \{F: F \text{ is symmetric about } 0, \int x^2 dF < \infty\}$ . Let  $\{U_i\}_{i=1}^{\infty}, \{V_i\}_{i=1}^{\infty}$  be two sequences of mutually independent identically distributed random variables with common distribution  $F$ , where  $F \in F$ . Let  $F_S(n)$  be based on  $\{U_i\}_{i=1}^{\infty}, \{V_i\}_{i=1}^{\infty}$ . Then for every  $a \geq 0$ ,

$$W(F_S(n), F, a) = 0.$$

Proof: Choose and fix  $\epsilon > 0$ . It is sufficient to find an  $F$  in  $F$  such that  $\lim_{n \rightarrow \infty} n^{-1} \log P_F(F_S(n) \geq a) \geq -\epsilon$ .

Let  $\alpha$  be chosen as in Theorem 6. Let  $\{X_i\}_{i=1}^{\infty}, \{Y_i\}_{i=1}^{\infty}$  be mutually independent identically distributed random variables such that  $X_1^2$  has a

Gamma  $(\alpha, 1)$  distribution. Let  $\{\delta_i\}_{i=1}^{\infty}$ ,  $\{\eta_i\}_{i=1}^{\infty}$  be mutually independent identically distributed symmetric Bernoulli random variables, independent of  $\{X_i\}_{i=1}^{\infty}$ ,  $\{Y_i\}_{i=1}^{\infty}$ . Let  $F_S(n)$  be based on  $\{U_i\}_{i=1}^{\infty} = \{\eta_i X_i\}_{i=1}^{\infty}$  and

$$\{V_i\}_{i=1}^{\infty} = \{\delta_i Y_i\}_{i=1}^{\infty}.$$

Let  $Z = 1 - (a + 1) \left( \sum_{i=1}^n Y_i^2 / \sum_{i=1}^n X_i^2 \right)$ . For any  $a > 0$ ,

$$P_F(F_S(n) \geq a) = P_F((a + 1) \left( \sum_{i=1}^n \delta_i Y_i \right)^2 - \left( \sum_{i=1}^n \eta_i X_i \right)^2 \geq n \left( \sum_{i=1}^n (a + 1) Y_i^2 - X_i^2 \right)) =$$

$$P_F\left(\left(\sum_{i=1}^n \eta_i X_i\right)^2 \leq n \left[ \sum_{i=1}^n (a + 1) Y_i^2 - \sum_{i=1}^n X_i^2 \right] + (a + 1) \left( \sum_{i=1}^n \delta_i Y_i \right)^2\right) \geq$$

$$P_F\left(\left(\sum_{i=1}^n \eta_i X_i\right)^2 \leq n \sum_{i=1}^n X_i^2 Z\right) = E_F P_F\left(\left(\sum_{i=1}^n \eta_i X_i\right)^2 \leq n \sum_{i=1}^n X_i^2 Z \mid \{X_i, Y_i\}_{i=1}^{\infty}\right) =$$

$$(3) \quad 1 - 2 E_F P_F\left(\left(\sum_{i=1}^n \eta_i X_i\right) / \left(\sum_{i=1}^n X_i^2\right)^{1/2} \geq Z^{1/2} \mid \{X_i, Y_i\}_{i=1}^{\infty}\right) \geq$$

$$(4) \quad E(1 - 2 \exp(-n I(\frac{1}{2}(1 + Z^{1/2}), \frac{1}{2}) \{X_i, Y_i\}_{i=1}^{\infty})) \geq$$

$$p(-n I(\frac{1}{2}(1 + Z^{1/2}), \frac{1}{2}) I(Z \geq 2/n) \mid \{X_i, Y_i\}_{i=1}^{\infty}) \geq$$

$$(1 - 2 \exp(-n I(\frac{1}{2}(1 + (2/n)^{1/2}), \frac{1}{2}))) P_F(Z \geq 2/n),$$

where (4) follows from (3) by Lemma 2.

By Lemma 1,  $1 - 2 \exp(-n I(\frac{1}{2}(1 + (2/n)^{1/2}), \frac{1}{2}))$  is asymptotic to  $(1 - 2 \exp(-1))$ .

By direct substitution,

$$P_F(Z \geq 2/n) = P_F\left(\sum_{i=1}^n X_i^2 / \sum_{i=1}^n Y_i^2 \geq (a + 1)n/(n - 2)\right).$$

Hence,  $\lim_{n \rightarrow \infty} n^{-1} \log P_F(F_S(n) \geq a) \geq \lim_{n \rightarrow \infty} n^{-1} \log P_F\left(\sum_{i=1}^n X_i^2 / \sum_{i=1}^n Y_i^2 \geq a + 1\right) = -\epsilon$ .

result follows.

One is tempted to conjecture the existence of a subclass of  $F$ , say  $F^*$ , such that  $W(F_S(n), F^*, a) > 0$ . No such class has been discovered, although a number of candidates have been ruled out. Notice that the construction in Theorem 7 is scale invariant. By suitable scaling, this construction can be made to satisfy any condition on the absolute first moment, the second moment, or the  $p$ th percentile of the underlying distribution. Restricting the underlying observations to the class of symmetric and bounded distributions does not yield a non-zero extremal large deviation rate, as can be seen below.

Theorem 8: Let  $H = \{F: F \text{ is symmetric about zero, and has support in } [-1, +1]\}$ . Let  $\{X_i\}_{i=1}^\infty, \{Y_i\}_{i=1}^\infty$  be two sequences of mutually independent identically distributed random variables with common distribution  $F$ , where  $F \in H$ . Let  $F_S(n)$  be based on  $\{X_i, Y_i\}_{i=1}^\infty$ . Then for every  $a \geq 0$ ,  $W(F_S(n), H, a) = 0$ .

Proof: Proceeding as before, let  $\epsilon > 0$  be chosen and fixed.

Let  $H$  be the distribution that assigns mass  $e^{-\epsilon}$  to 0 and is uniformly distributed over  $(-1, +1)$  with probability  $(1 - e^{-\epsilon})$ . Then for every  $a \geq 0$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} n^{-1} \log P_H(F_S(n) \geq a) &\geq \\ \lim_{n \rightarrow \infty} n^{-1} \log P_H(F_S(n) = +\infty) &= \\ \lim_{n \rightarrow \infty} n^{-1} \log P_H(s_{x,n}^2 > 0, s_{y,n}^2 = 0) &= \dots \\ \lim_{n \rightarrow \infty} n^{-1} \{ \log P(s_{x,n}^2 \neq 0) + \log P(s_{y,n}^2 = 0) \} &= \\ \lim_{n \rightarrow \infty} n^{-1} \{ \log(1 - e^{-n\epsilon}) + \log(e^{-n\epsilon}) \} &= \\ -\epsilon. \quad \square \end{aligned}$$

#### IV. Efficiencies of Tests for Equality of Variances

Consider the problem of testing for equality of variances of two populations, with underlying distributions unspecified. The usual procedure is to reject the null hypothesis, equality of variances, if the statistic  $F_s(n)$  exceeds some specified value. Call this test  $T_1$ . Let us consider a competitor to  $T_1$ .

Moses (1963) has proposed a distribution-free test for equality of variances. Moses recommends randomly allocating  $\{X_i\}_{i=1}^{kn}$  into  $n$  subsamples of size  $k$ , and calculating sample variances. This is repeated with  $\{Y_i\}_{i=1}^{kn}$ , and the two collections of sample variances are then compared by means of the Wilcoxon rank sum test. Call this test  $T_2$ .

It is known that the half slope of the Wilcoxon rank sum test is positive. Hence, the half slope of  $T_2$  is also positive. By Theorem 7, the half slope of  $T_1$  is zero at every alternative. Hence, at every alternative, the Bahadur efficiency of the F-test, relative to Moses' test, is zero.

While the t-test performs relatively well against distribution free competitors, the F-test performs as poorly as possible. One is tempted to propose a "t-test for variances" of the form

$$S_n = (s_{x,n}^2 - s_{y,n}^2) / (s_{x,n}^4 + s_{y,n}^4)^{1/2}.$$

The statistic  $S_n$  can be shown to be a monotone increasing function of  $F_s(n)$  if  $F_s(n) > 0$  and a monotone decreasing function of  $F_s(n)$  if  $F_s(n) < 0$ . Hence, the half slope of  $S_n$  is zero at every alternative, just as the half slope of  $F_s(n)$  is.

Table 1: Large Deviation Rates of the F-statistic

Deviation a	Normal Observations $-\log(2(a+1)^{\frac{1}{2}}/(a+2))$	Bernoulli Observations $I(\frac{1}{2}(1 + (a/a+1)^{\frac{1}{2}}), \frac{1}{2})$
.05	.00030	.02400
.10	.00114	.04617
.15	.00244	.06672
.20	.00415	.08582
.25	.00621	.10363
.30	.00858	.12029
.35	.01122	.13591
.40	.01652	.15058
.45	.01716	.16440
.50	.02041	.17744
.55	.02382	.18977
.60	.02736	.20144
.65	.03102	.21251
.70	.03479	.22304
.75	.03865	.22305
.80	.04257	.24259
.85	.04658	.25169
.90	.05064	.26038
.95	.05474	.26869
1.00	.05889	.27665
1.25	.08004	.31188
2.50	.10147	.34102
2.75	.12281	.36557
3.00	.14384	.38658



Table 1  
(continued)

Deviation	Normal Observations	Bernoulli Observations
$a$	$-\log(2(a+1)^{1/2}/(a+2))$	$I(\frac{1}{2}(1+(a/a+1)^{1/2}), \frac{1}{2})$
3.50	.18455	.42072
4.00	.22314	.44737
5.00	.29389	.48651
6.00	.35688	.51406
7.00	.41334	.53456
8.00	.46436	.55052
9.00	.51083	.56332
10.00	.56358	.57618
11.00	.59281	.58264
12.00	.62934	.59014
13.00	.66344	.59660
14.00	.69537	.60225
15.00	.72542	.60721
20.00	.85351	.62526
$\infty$	$\infty$	$\log 2$

## REFERENCES

- Bahadur, R. R. (1960). Stochastic comparison of tests. Annals of Mathematical Statistics 31, 276-295.
- \_\_\_\_\_. (1967). Rates of convergence of estimates and test statistics. Annals of Mathematical Statistics 38, 303-325.
- \_\_\_\_\_. (1971). Some Limit Theorems in Statistics, SIAM, Philadelphia.
- Chernoff, H. (1952). A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. Annals of Mathematical Statistics 23, 493-507.
- Jones, D. H. and Sethuraman, J. (1978). The efficiency of some nonparametric competitors to the t-test. Annals of Mathematical Statistics 27, 324-335.
- Klotz, J. (1965). Alternative efficiencies for signed rank tests. Annals of Mathematical Statistics 36, 1759-1766.
- Hodges, J. L. and Lehmann, E. L. (1956). The efficiency of some nonparametric competitors to the t-test. Annals of Mathematical Statistics 27, 324-335.
- Moses, L. E. (1963). Rank tests for dispersion. Annals of Mathematical Statistics 34, 973-983.
- Pitman, E. J. G. (1949). Lecture notes on nonparametric statistical inference. Columbia University.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU No. M529 USARO No. D-42	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE THE EXTREMAL LARGE DEVIATION RATE OF THE F-STATISTIC	5. TYPE OF REPORT & PERIOD COVERED Technical Report	6. PERFORMING ORG. REPORT NUMBER FSU Statistics Report M529
7. AUTHOR(S) L. Barker J. Sethuraman	8. CONTRACT OR GRANT NUMBER(S) USARO DAA29-76-G00278 <sup>eu</sup>	9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
10. PERFORMING ORGANIZATION NAME & ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	11. CONTROLLING OFFICE NAME & ADDRESS U.S. Army Research Office-Durham P.O. Box 12211 Research Triangle Park, NC 27709	12. REPORT DATE October, 1979
13. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	14. NUMBER OF PAGES 15	15. SECURITY CLASS(of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this report) Approved for public release; distribution unlimited.	17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)	18. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. SUPPLEMENTARY NOTES		
20. KEY WORDS Bahadur efficiency, extremal large deviations, F-statistic, nonparametric tests.		
21. ABSTRACT The extremal large deviation rate of the F-statistic over a large class is determined. It is shown that the Bahadur efficiency of the F-test relative to Moses' rank test for equality of variances is zero at every alternative when the distribution of the underlying observations is unspecified.		